

---

# Phaeodactylum metabolism converges to phosphoenolpyruvate formation during growth under different light conditions

Parisa Heydarizadeh<sup>1</sup>, Wafâa Boureba<sup>1</sup>, Brigitte Moreau<sup>1</sup>, Bing Huang<sup>1</sup>, Ewa Lukomska<sup>2</sup>, Véronique Martin-Jézéquel<sup>3</sup>, Gaëtane Wielgosz-Collin<sup>4</sup>, Aurélie Couzinet-Mossion<sup>4</sup>, Justine Marchand<sup>1</sup>, and Benoît Schoefs<sup>\*†1</sup>

<sup>1</sup>MicroMar, Mer Molécules Santé (MMS) – University of Le Mans, IUML FR 3473 CNRS, – 72000 Le Mans, France

<sup>2</sup>Physiologie et biotechnologie des Algues (PBA) – Institut Français de Recherche pour l'Exploitation de la Mer (IFREMER) – Rue de l'Île d'Yeu BP 21105 Nantes cedex 3, France

<sup>3</sup>Littoral, Environnement et Sociétés (LIENSs) – CNRS : UMR7266 – Université de La Rochelle, 2 rue Olympe de Gouge, 17042 La Rochelle Cedex 1, France

<sup>4</sup>ChimiMar, Mer Molécules Santé (MMS) – University of Nantes, IUML FR 3473 CNRS, – 44035 Nantes, France

## Résumé

Gigatons of CO<sub>2</sub> are yearly converted into organic compounds by photosynthetic organisms and diatoms are among the most important contributors in this process. Both the capacity of carbon fixation and the fate of fixed carbon are strongly impacted by the environmental factors. Because light can constitute a stressful factor when present in inadequate amount, various mechanisms have been developed by diatoms to adapt very efficiently to changing irradiance. The molecular mechanisms on which light acclimation in diatoms relies remain largely unknown. To obtain further information on these mechanisms, the impacts of different growth photon flux densities (30 (LL), 300 (ML), 1000 (HL)  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ) were studied at the physiological, biochemical and molecular levels in the model diatom *Phaeodactylum tricorutum*. The integrated results indicate that the impact of ML and HL on diatom cells were similar but quite different from LL. In addition of light and growth phase could affect pigments concentration and primary metabolites. The trend of gene expression coding enzymes in central carbon metabolism pathways differed under LL, but the orientation of the metabolisms was toward either phosphoenolpyruvate (PEP) or pyruvate formation under all three light intensities. LL provided a condition for cells to accumulate chrysolaminarin and lipids, while ML mostly stimulated lipid synthesis. A significant increase in the amount of proteins was observed under HL. We concluded that PEP and pyruvate are key intermediates in diatom to synthesis valuable compounds such as lipids, proteins, aromatic amino acids and polyphenolics.

**Mots-Clés:** carbon metabolism, stress physiology, diatom, light stress, metabolic reorientation

---

\*Intervenant

†Auteur correspondant: benoit.schoefs@univ-lemans.fr